

Assessment of vegetation indices derived from UAV images for predicting biometric variables in bean during ripening stage

Assessment of vegetation indices derived from UAV images for predicting biometric variables in bean during ripening stage

Javier Quille-Mamani^{1*}, Rossana Porras-Jorge¹, David Saravia-Navarro^{1, 2}, Jordán Herrera¹, Julio Chávez-Galarza¹, Carlos I. Arbizu¹, Lamberto Valqui-Valqui¹

ABSTRACT

Here, we report the prediction of vegetative stages variables of canary bean crop employing RGB and multispectral images obtained from UAV during the ripening stage, correlating the vegetation indices with biometric variables measured manually in the field. Results indicated a highly significant correlation of plant height with eight vegetation indices derived from UAV images from the canary bean, which were evaluated by multiple regression models, obtaining a maximum correlation of $R^2 = 0.79$. On the other hand, the estimated indices of multispectral images did not show significant correlations.

Keywords: Vegetation indices; precision agriculture; RGB images.

RESUMEN

Here, we report the prediction of vegetative stages variables of canary bean crop employing RGB and multispectral images obtained from UAV during the ripening stage, correlating the vegetation indices with biometric variables measured manually in the field. Results indicated a highly significant correlation of plant height with eight vegetation indices derived from UAV images from the canary bean, which were evaluated by multiple regression models, obtaining a maximum correlation of $R^2 = 0.79$. On the other hand, the estimated indices of multispectral images did not show significant correlations.

Palabras clave: Vegetation indices; precision agriculture; RGB images.

Introduction

Bean (*Phaseolus vulgaris* L.) is an essential legume as a source of protein and dietary fiber for millions of people (Los *et al.*, 2018). It is widely cultivated for its enormous genetic diversity. It is a nitrogen-fixing plant, highly adaptable and productive in a wide range of environments (Shamseldin & Velázquez, 2020). This crop will likely play a key role in guaranteeing food security for millions of people worldwide in the near future (Mecha *et al.*, 2018). In Peru in 2019, a total of 73,298 ha of beans were cultivated, representing 0.8% of the Gross Value of Agricultural Production (GVaP) (MIDAGRI, 2020). Canary bean is the most outstanding cultivar on the Peruvian coast due to its

preference in the national diet and cultural aspects. Therefore, due to the importance of this crop, it is a great challenge to monitor its development together with appropriate agronomic management in the field (Shakoor *et al.*, 2017) in the context of climate change. Quantitative evaluations of biometric variables such as plant height, leaf area index, and chlorophyll content influence yield and become a high priority under precision agriculture (Eggen *et al.*, 2019). Efficient and non-destructive crop growth monitoring is essential for appropriate crop management and key for digital agriculture (Wang *et al.*, 2019). Determining data manually requires a significant amount of time and resources (measuring equipment, reagents, and researchers, among others). Important technological advances

¹ Dirección de Desarrollo Tecnológico Agrario, Instituto Nacional de Innovación Agraria (INIA), Av. La Molina 1981, Lima 15024, Lima, Perú.

² Facultad de Agronomía, Universidad Nacional Agraria La Molina, Av. La Molina s/n, Lima 15024, Lima, Perú.

* Corresponding author: (geomatica@inia.gob.pe)

have been implemented, such as the use of unmanned aerial vehicles to increase agricultural production with limited resources (UAVs) (Khan *et al.*, 2021).

UAVs are tools that provide new alternatives for monitoring crops without direct contact with them (Qi *et al.*, 2020), allowing the prediction of crop development in a spatial-temporal way. The sensors coupled to a UAV allow estimating vegetative development variables with different vegetative indices (Gano *et al.*, 2021). Recently, in the north of China, (Guo *et al.*, 2020) used UAV images to predict yield in corn. Similarly, other researchers predicted aerial biomass in various crops such as sunflower and wheat (Du & Noguchi, 2017; Vega *et al.*, 2015). In Kenya, RGB images were used to estimate cassava's growth and nutritional yield (Wasonga *et al.*, 2021). Since evaluations are necessary during the development of crops and as beans are one of the main foods in the basic family basket, research must be conducted to guarantee food security under the context of climate change. Therefore, the objective of the present work was to estimate the prediction of plant height, chlorophyll content, and leaf area index through multispectral images obtained from UAV of canary bean crops during the ripening stage.

Materials and methods

Study site

The study site is located in the research field of the National Institute of Agrarian Innovation (INIA for its acronym in Spanish) (12°4'30.10''S, 76°56'33.86''W, 240 m.a.s.l) (Figure 1). The

experiment was developed during the winter-spring seasons (Jun 26-Oct 2020), using commercial canary beans. An experimental field of 0.30 ha (Figure 1) with distances between plants and rows of 0.2 m and 0.9 m, respectively, was used for this research. We used drip irrigation with a distance between drippers of 0.2 m with a flow of 1.20 l h⁻¹. The study site is arid, with an average temperature of 19.2 °C and a total annual rainfall of 8 mm.

Field data collection

Three biometric variables were recorded on 16 plots of 7 m x 5 m at 90, 97, and 101 days after sowing (DAS). Plant height (cm) was measured manually from the soil surface to the highest stem apex. Leaf area index (LAI) was estimated from digital images taken on the canopy cover of the plant, as indicated by (Peng *et al.*, 2021). was with the CCM-300 equipment (OptiSciences Hudson, NH, USA) measured Chlorophyll content (mg m⁻²) following the methodology proposed by (Guo *et al.*, 2020).

UAV RGB and multispectral image acquisition and processing

We used a Quadcopter-type UAV platform, DJI Phantom 4 Pro (Shenzhen Dajiang Baiwang Technology Co., Ltd., Shenzhen, China), with a built-in 4864 × 3648 pixel resolution RGB camera. In addition, a multispectral sensor Parrot Sequoia (Parrot SA, France), which is a synchronized array of 4 single-band multispectral cameras with 1.2 MP global shutter attached, taking images in green




Figure 1. Location of the 64 canary bean research plots. The ground control points (GCP) are shown, and the 16 plots sampled for the estimation of vegetation indices are shown in blue rectangles.

(550 nm), red (660 nm), red edge (735 nm) and near infrared (790 nm). The images were collected at 90, 97, and 101 DAS during sunny days with wind speeds lower than 12 m s⁻¹, from 11:00 to 13:00 hours approximately on the 16 research plots, as detailed in Figure 1. The flight plan was established with the Pix4Dcapture software (v. 4.12.1, Pix4D SA, Prilly, Switzerland), considering a frontal and lateral overlap of 80%, the height of 30 m, speed of 2.8 m s⁻¹, and the camera focused at nadir position (perpendicular to the ground surface), allowing to obtain a resolution of 0.8 cm and 2.5 cm for RGB and multispectral images, respectively. Figure 2 shows a flow chart for data acquisition from the UAV platform to image processing. Image processing was performed with Pix4Dmapper Pro software (v. 4.3.33, Pix4D SA, Prilly, Switzerland), according to the following steps: (i) alignment of geolocated images, (ii) generation of point clouds and geometric correction, and (iii) generation of the digital surface and orthomosaic model using the inverse distance weighting method. The geometric correction was performed considering nine ground

control points (GCP) (Figure 1) previously installed and registered with differential GNSS (Global Navigation Satellite System) (South Galaxy G1 model, South Surveying & Mapping Instrument Co. Ltd, Guangdong, China).

Calculation of vegetation indices

The vegetation indices were calculated from RGB images and were carried out prior to bands normalization ($R = \text{red}$, $G = \text{green}$, and $B = \text{blue}$) in the Pix4Dmapper software. In addition, the multispectral bands were obtained ($R = \text{red}$, $G = \text{green}$, $RE = \text{red border}$, and $NIR = \text{near infrared}$). In this research, 19 vegetation indices were estimated to assess biometric parameters (Table 1).

Statistical analysis: Biometric variables (plant height, leaf area index, and chlorophyll content) and vegetation indices were correlated using the following statistical indicators: Pearson's correlation coefficient (r) and coefficient of determination (R^2); both were calculated with R software v. 4.1.0 (R Core Team, 2021).




Figure 2. Flowchart for the acquisition and processing of images for the calculation of the vegetation indices of a canary bean crop.

Table 1. Vegetation index calculations from Red-Green-Blue and Multispectral images.

Abbrev	Definition	type	Reference
NGRDI	$(G - R)/(G + R)$		(Han <i>et al.</i> , 2019)
ExG	$2xG - R - B$		(Han <i>et al.</i> , 2019)
ExR	$1.4xR - G$		(Bendig <i>et al.</i> , 2015)
ExB	$1.4xB - G$		(Ballesteros <i>et al.</i> , 2018; Guo <i>et al.</i> , 2020)
IKAW	$(R - B)/(R + B)$		(Du & Noguchi, 2017)
GRR1	G/R		(Beniaich <i>et al.</i> , 2019)
ExGR	$ExG - ExR$		(Lussem <i>et al.</i> , 2018)
GBDI	$G - B$	RGB	(Bendig <i>et al.</i> , 2015; Lussem <i>et al.</i> , 2018)
MGRVI	$(G \times G - R \times R)/(G \times G + R \times R)$		(Wan <i>et al.</i> , 2018)
RGBVI	$(G \times G - B \times R)/(G \times G + B \times R)$		(Lussem <i>et al.</i> , 2018)
VDVI	$(2 \times G - R - B)/(2 \times G + R + B)$		(Wan <i>et al.</i> , 2018)
VARI	$(G - R)/(G + R - B)$		(Beniaich <i>et al.</i> , 2019)
RGRI	R/G		
NGBDI	$(G - B)/(G + B)$		
CIVE	$(0.441 \times R - 0.8818 \times G + 0.385 \times B + 18.787)$		
NDVI	$(NIR - r)/(NIR + r)$		(Ranjan <i>et al.</i> , 2019; Wasonga <i>et al.</i> , 2021)
GNDVI	$(NIR - g)/(NIR + g)$	Multispectral	(Sankaran <i>et al.</i> , 2019)
SAVI	$[(NIR - r)/(NIR + r + L)](1 + L)$		(Ranjan <i>et al.</i> , 2019)
NDREI	$(NIR - re)/(NIR + re)$		(Hassan <i>et al.</i> , 2018; Ranjan <i>et al.</i> , 2019)

Results and discussion

Correlation between spectral indices and growth variables

The correlation analysis between the vegetation indices and growth variables was carried out for 90, 97, and 101 DAS, finding significant correlations (p -value < 0.05) at 97 DAS for plant height. According to (Schirrmann *et al.*, 2016), the plant height and coverage derived from the UAV images became important because chlorophyll content decreases in the wheat leaves with further propagation of senescence. In contrast, the other variables (chlorophyll content and leaf area index) did not register significant correlations, so we decided not to consider them for the multiple linear regression model. Plant height is an important variable since it depends on the growth rate of crops (Rai *et al.*, 2020). In Table 2, 19 vegetation indices correlated to plant height are detailed. Four indices (RGBVI, VDVI, ExGR, and ExB) presented high significance (p -value < 0.01), eight indices (ExR, GBDI, NGBDI, NGRDI, MGRVI, VARI, CRRI, and RGRI) showed significance (p -value < 0.05), and four indices (ExB, ExR, IKAW, and ExG) were inversely proportional. No multispectral index presented significant correlations (NDVI, SAVI, GNDVI, and NDREI).

Table 2. Vegetation indices correlated with plant height recorded at 97 DAS.

Indice	plant height	
	<i>r</i> Pearson	<i>p</i> -value
NDVI	0.39	0.135 ns
SAVI	0.40	0.125 ns
GNDVI	0.33	0.215 ns
NDREI	0.37	0.156 ns
CIVE	0.22	0.418 ns
RGBVI	0.64	0.008 **
ExR	-0.62	0.011 *
GBDI	0.63	0.010 *
VDVI	0.64	0.007 **
NGBDI	0.60	0.014 *
NGRDI	0.61	0.011 *
MGRVI	0.61	0.012 *
IKAW	-0.18	0.502 ns
VARI	0.59	0.017 *
ExGR	0.63	0.009 **
ExG	-0.32	0.223 ns
CRRI	0.62	0.010 *
RGRI	0.60	0.014 *
ExB	-0.66	0.005 **

**Altamente significativos, * Significativo, ns: No significativo.

Our results agree with those of (Hassan *et al.*, 2018), who stated that the vegetation indices decrease after flowering in wheat, causing fluctuations during senescence.

Prediction model for estimating plant height.

Multiple linear regression analyzes were performed for three prediction models after correlating the vegetation indices and plant height at 97 DAS, as detailed in Table 3. Figure 3 shows the three prediction models for the measured plant height and its predicted values: (i) for model I, 10 vegetation indices were used (RGBVI, GBDI, VDVI, NGBDI, NGRDI, MGRVI, VARI, ExGR, CRRI, RGRI) with significant correlations $0.59 < r < 0.64$ (p -value < 0.05), generating a predictive model with an $R^2 = 0.79$ (p -value < 0.01), (ii) in model II four vegetation indices were considered (RGBVI, GBDI, VDVI, ExGR) with significant correlations of $0.63 < r$

< 0.64 , (p -value < 0.01), obtaining a predictive model with $R^2 = 0.60$, and (iii) to generate the prediction model III, three vegetation indices were considered (ExR, ExG, ExB) with negative correlations $-0.66 < r < -0.32$, generating a predictive model with an $R^2 = 0.46$. Although for model II, four vegetation indices with high correlation values were chosen, their prediction decreases because it presents an R^2 lower than model I but higher than that of model III. As we add a greater number of indices to the predictive model, it is more reliable. However, model II with four indices also predicts high significance; we would use it for data analysis if we had fewer resources. (Bendig *et al.*, 2015) reported plant height estimation with two vegetative indices (GRVI and

Table 3. Multivariate linear regression models between vegetation indices and plant height recorded during ripening stage for canary bean.

Prediction model	Spectral Index	Regression Coefficient	Intercept	Model R ²
I	RGBVI	-2893.1	5934.9	0.79
	GBDI	21228.7		
	VDVI	5500.7		
	NGBDI	-8715.3		
	NGRDI	5934.9		
	MGRVI	-9715.4		
	VARI	3526.0		
	EXGR	-5071.7		
	CRRI	-139.1		
	RGRI	1657.0		
II	RGBVI	-3556	2709	0.60
	GBDI	4832		
	VDVI	-1332		
	EXGR	-3179		
III	EXR	-91.31	51.45	0.46
	EXG	210		
	EXB	-216.05		




Figure 3. Correlation between the estimated data and plant height recorded for three models, A) Model I is based on ten vegetation indices, B) Model II is based on four vegetation indices, and C) Model III is based on three vegetation indices with inverse correlation.

RGBVI) evaluated during different stages of barley through the use of multiple linear regression models.

To our best knowledge, this is the first research work in Peru using technology such as UAV for agronomic monitoring of canary beans. A highly significant correlation for plant height was obtained from eight estimated vegetation indices from RGB images in this crop at the ripening stage. It was possible to predict the size of this crop with high performance ($R^2 = 0.79$) using model I. Recent studies (Parker *et al.*, 2020; Wasonga *et al.*, 2021) reported similar results for beans and cassava crops, demonstrating the utility of UAVs and vegetation indices to compile biometric variables.

Conclusions

A highly significant correlation was obtained for plant height with eight RGB image vegetation indices in the commercial canary bean crop. The obtained indices from multispectral images did not show significant correlations. The predictive model I with $R^2 = 0.79$ predicts plant height with good accuracy. The use of these tools is important due to the need to conduct more agronomic evaluations promptly in crops and mainly in those that are cultivated in arid environments such as the Peruvian

coast. We are currently conducting experiments with new technologies using remote sensors such as UAV to predict biometric variables estimated from vegetative indices obtained from RGB and multispectral images in bean crops and others of national importance. In addition, images from the PeruSat-1 (Peru) and Kompsat 3 (South Korea) satellites will be used, allowing timely monitoring and prediction of crop development under different agronomic management scenarios (irrigation, fertilization, among others) on a larger scale.

Acknowledgements

We are grateful to PIP 2281955, "Installation of the agricultural technology research service specialized in climate change for the agricultural sector" and PIP 2449640, "Creation of the precision agriculture service in the departments of Lambayeque, Huancavelica, Ucayali and San Martín 4 Departments".

We are grateful to PIPs 2281955 and 2449640 for providing funding. Likewise, thanks to the students of the Agrotechnology Research Circle (CIATEC for its acronym in Spanish) of the Faculty of Agronomy, La Molina National Agrarian University, for their support during field evaluations.

Literature cited

- Ballesteros, R.; Ortega, J.F., Hernandez, D.; Moreno, M.A. 2018. Onion biomass monitoring using UAV-based RGB imaging. *Precision Agriculture*, 19 (5): 840-857.
- Bendig, J.; Yu, K.; Aasen, H.; Bolten, A.; Bennertz, S.; Broscheit, J.; Gnyp, M.L.; Bareth, G. 2015. Combining UAV-based plant height from crop surface models, visible, and near infrared vegetation indices for biomass monitoring in barley. *International Journal of Applied Earth Observation and Geoinformation*, 39: 79-87.
- Beniaich, A.; Naves Silva, M.L.; Avalos, F.A.P.; Menezes, M.D.; Candido, B.M. 2019. Determination of vegetation cover index under different soil management systems of cover plants by using an unmanned aerial vehicle with an onboard digital photographic camera. *Semina-Ciencias Agrarias*, 40 (1): 49-66.
- Du, M., Noguchi, N. 2017. Monitoring of Wheat Growth Status and Mapping of Wheat Yield's within-Field Spatial Variations Using Color Images Acquired from UAV-camera System. *Remote Sensing*, 9 (3): 289.
- EGgen, M.; Ozdogan, M.; Zaitchik, B.; Ademe, D.; Foltz, J.; Simane, B. 2019. Vulnerability of sorghum production to extreme, sub-seasonal weather under climate change. *Environmental Research Letters*, 14 (4): 045005.
- Gano, B.; Dembele, J.S.B.; Ndour, A.; Luquet, D.; Beurier, G.; Diouf, D.; Audebert, A. 2021. Using UAV Borne, Multi-Spectral Imaging for the Field Phenotyping of Shoot Biomass, Leaf Area Index and Height of West African Sorghum Varieties under Two Contrasted Water Conditions. *Agronomy*, 11 (5): 850.
- Guo, Y.; Wang, H.; Wu, Z.; Wang, S.; Sun, H., Senthilnath, J.; Wang, J.; Robin Bryant, C.; Fu, Y. 2020. Modified Red Blue Vegetation Index for Chlorophyll Estimation and Yield Prediction of Maize from Visible Images Captured by UAV. *Sensors*, 20 (18): 5055.
- Han, L.; Yang, G.; Dai, H.; Xu, B.; Yang, H.; Feng, H.; Li, Z.; Yang, X. 2019. Modeling maize above-ground biomass based on machine learning approaches using UAV remote-sensing data. *Plant Methods*, 15 (1): 10.
- Hassan, M. A.; Yang, M.; Rasheed, A.; Jin, X.; Xia, X.; Xiao, Y.; He, Z. 2018. Time-Series Multispectral Indices from Unmanned Aerial Vehicle Imagery Reveal Senescence Rate in Bread Wheat. *Remote Sensing*, 10 (6): 809.
- Khan, N.; Ray, R.L.; Sargani, G.R.; Ihtisham, M.; Khayyam, M.; Ismail, S.

2021. Current Progress and Future Prospects of Agriculture Technology: Gateway to Sustainable Agriculture. *Sustainability*, 13 (9): 4883.
- Los, F.G.B.; Zielinski, A.A.F.; Wojeicchowski, J.P.; Nogueira, A.; Demiate, I.M.
2018. Beans (*Phaseolus vulgaris* L.): Whole seeds with complex chemical composition. *Current Opinion in Food Science*, 19, 63-71.
- Lussem, U.; Bolten, A.; Gnyp, M.L.; Jasper, J.; Bareth, B.
2018. Evaluation of RGB-based vegetation indices from UAV imagery to estimate forage yield in grassland. *Int. Arch. Photogramm. Remote Sens. Spatial Inf. Sci.*, 42 (3): 1215-1219.
- Mecha, E.; Figueira, M.E.; Bronze, M.C.V.P.; M. do R.
2018. Two Sides of the Same Coin: The Impact of Grain Legumes on Human Health: Common Bean (*Phaseolus vulgaris* L.) as a Case Study. In *Legume Seed Nutraceutical Research*. DOI: 10.5772/intechopen.78737
- MIDAGRI, D.
2020. Plan Nacional de Cultivos: Campaña Agrícola 2019-2020. Ministerio de Agricultura. Lima, Perú. 322 p.
- Parker, T.A.; Palkovic, A.; Gepts, P.
2020. Determining the Genetic Control of Common Bean Early-Growth Rate Using Unmanned Aerial Vehicles. *Remote Sensing*, 12 (11): 1748.
- Peng, X.; Han, W.; Ao, J.; Wang, Y.
2021. Assimilation of LAI Derived from UAV Multispectral Data into the SAFY Model to Estimate Maize Yield. *Remote Sensing*, 13 (6): 1094.
- Qi, H.; Zhu, B.; Wu, Z.; Liang, Y.; Li, J.; Wang, L.; Chen, T.; Lan, Y.; Zhang, L.
2020. Estimation of Peanut Leaf Area Index from Unmanned Aerial Vehicle Multispectral Images. *Sensors*, 20 (23): 6732.
- R Core Team.
2021. *RStudio: Integrated Development Environment para R*, Boston, MA: RStudio, Inc.
- Rai, A.; Sharma, V.; Heitholt, J.
2020. Dry Bean [*Phaseolus vulgaris* L.] Growth and Yield Response to Variable Irrigation in the Arid to Semi-Arid Climate. *Sustainability*, 12 (9): 3851.
- Ranjan, R.; Chandel, A.K.; Khot, L.R.; Bahlol, H.Y.; Zhou, J.; Boydston, R.A.; Miklas, P.N.
2019. Irrigated pinto bean crop stress and yield assessment using ground based low altitude remote sensing technology. *Information Processing in Agriculture*, 6 (4): 502-514.
- Sankaran, S.; Quirós, J.J.; Miklas, P.N.
2019. Unmanned aerial system and satellite-based high resolution imagery for high-throughput phenotyping in dry bean. *Computers and Electronics in Agriculture*, 165: 104965.
- Schirrmann, M.; Giebel, A.; Gleiniger, F.; Pflanz, M.; Lentschke, J.; Dammer, K.-H.
2016. Monitoring Agronomic Parameters of Winter Wheat Crops with Low-Cost UAV Imagery. *Remote Sensing*, 8 (9): 706.
- Shakoor, N.; Lee, S.; Mockler, T.C.
2017. High throughput phenotyping to accelerate crop breeding and monitoring of diseases in the field. *Current Opinion in Plant Biology*, 38: 184-192.
- Shamseldin, A.; Velázquez, E.
2020. The promiscuity of *Phaseolus vulgaris* L. (common bean) for nodulation with rhizobia: A review. *World Journal of Microbiology and Biotechnology*, 36 (5): 1-12.
- Vega, F.A.; Ramírez, F.C.; Saiz, M.P.; Rosúa, F.O.
2015. Multi-temporal imaging using an unmanned aerial vehicle for monitoring a sunflower crop. *Biosystems Engineering*, 132: 19-27.
- Wan, L.; Li, Y.; Cen, H.; Zhu, J.; Yin, W.; Wu, W.; Zhu, H.; Sun, D.; Zhou W.; He, Y.
2018. Combining UAV-Based Vegetation Indices and Image Classification to Estimate Flower Number in Oilseed Rape. *Remote Sensing*, 10 (9): 1484.
- Wang, Y.; Zhang, K.; Tang, C.; Cao, Q.; Tian, Y.; Zhu, Y.; Cao, W.; Liu, X.
2019. Estimation of Rice Growth Parameters Based on Linear Mixed-Effect Model Using Multispectral Images from Fixed-Wing Unmanned Aerial Vehicles. *Remote Sensing*, 11 (11): 1371.
- Wasonga, D.O.; Yaw, A.; Kleemola, J.; Alakukku, L.; Mäkelä, P.S.A.
2021. Red-Green-Blue and Multispectral Imaging as Potential Tools for Estimating Growth and Nutritional Performance of Cassava under Deficit Irrigation and Potassium Fertigation. *Remote Sensing*, 13 (4): 598.